

ATOMIC NUMBER DENSITY

Number of Atoms (n) and Number Density (N)

The number of atoms or molecules (n) in a mass (m) of a pure material having atomic or molecular weight (M) is easily computed from the following equation using Avogadro's number ($N_{Av} = 6.022 \times 10^{23}$ atoms or molecules per gram-mole):

$$n = \frac{m N_{Av}}{M} \quad (1)$$

In some situations, the *atomic number density* (N), which is the concentration of atoms or molecules per unit volume (V), is an easier quantity to find when the density (ρ) is given

$$N = \frac{n}{V} = \frac{\rho N_{Av}}{M} \quad (2)$$

Number Density for Compounds

For a chemical compound (mixture) Z , which is composed of elements X and Y , the number (atom) density of the compound is calculated from

$$N_Z = N_{mix} = \frac{\rho_{mix} N_{Av}}{M_{mix}} \quad (3)$$

In some cases, the desired quantity is the number density of the compound constituents. Specifically, if $Z = X_p Y_q$, then there are p atoms of X and q atoms of Y for every molecule of Z ; hence

$$\begin{aligned} N_X &= p N_Z \\ N_Y &= q N_Z \end{aligned} \quad (4)$$

Example: Calculate the number density of natural uranium in UO_2 with $\rho_{UO_2} = 10.5 \text{ g/cm}^3$.

$$N_U = N_{UO_2} = \frac{\rho_{UO_2} N_{Av}}{M_{UO_2}} = \frac{(10.5 \text{ g/cm}^3)(6.022 \times 10^{23} \text{ atoms/mol})}{[238.0289 + 2(15.9994)] \text{ g/mol}} = 2.34 \times 10^{22} \text{ atoms/cm}^3$$

Number Density Given Atom Fraction (Abundance)

Oftentimes, it is necessary to compute the concentration of an individual isotope j given its fractional presence (abundance) γ_j in the element

$$\gamma_j = \frac{\text{Number of atoms of isotope } j}{\text{Total number of atoms of the element}} \quad (5)$$

Many times, the fraction γ_j is stated as an atom percent, which is abbreviated a/o. The atomic number density of isotope j is then

$$N_j = \gamma_j N_{elem} = \frac{\gamma_j \rho_{elem} N_{Av}}{M_{elem}} \quad (6)$$

If the element has a non-natural abundance of its isotopes (that is, the elemental material is either *enriched* or *depleted*), then it is necessary to compute the atomic weight of the element (M_{elem}) from the sum of all the atomic weights of the isotopes (M_j) rather than use the tabulated M_{elem} value found in a reference

$$M_{elem} = \sum \gamma_j M_j \quad (7)$$

Example: Find the U-235 concentration for 3 a/o in UO₂.

Solution: To solve this example, Equations (4), (3) and (7) are progressively substituted into Eq. (6).

$$\begin{aligned}
 N_{\text{U-235}} &= \gamma_{\text{U-235}} N_{\text{U}} = \gamma_{\text{U-235}} N_{\text{UO}_2} \\
 &= \gamma_{\text{U-235}} \frac{\rho_{\text{UO}_2} N_{\text{Av}}}{M_{\text{UO}_2}} = \frac{\gamma_{\text{U-235}} \rho_{\text{UO}_2} N_{\text{Av}}}{\gamma_{238} M_{238} + \gamma_{235} M_{235} + 2 M_{\text{O}}} \\
 &= \left(0.03 \frac{\text{atoms-}^{235}\text{U}}{\text{atoms - U}} \right) \frac{(10.5 \text{ g/cm}^3)(6.022 \times 10^{23} \text{ atoms/mol})}{[(238.05)(0.97) + (235.04)(0.03) + 2(15.9994)] \text{ g/mol}} \\
 &= 7.03 \times 10^{20} \text{ atoms/cm}^3
 \end{aligned}$$

Number Density Given Weight Fraction (Enrichment)

Other times, when working with nuclear fuels such as uranium, the *enrichment* may be specified in terms of weight percent or weight fraction, ω_i , of isotope i :

$$\omega_i = \frac{\text{Mass of isotope } i}{\text{Total mass of the element}} \quad (8)$$

The atomic number density of isotope i is

$$N_i = \frac{\rho_i N_{\text{Av}}}{M_i} = \frac{\omega_i \rho_{\text{elem}} N_{\text{Av}}}{M_i} \quad (9)$$

Clearly, if the material is enriched, then the atomic weight of the material differs from its natural reference value, and the enriched atomic weight, if needed, should be computed from

$$\frac{1}{M_{\text{elem}}} = \sum_i \frac{\omega_i}{M_i} \quad (10)$$

Example: Find the U-235 concentration for 4% enriched UO₂.

Solution: First, compute the molecular weight of the enriched uranium, which is basically 4% U-235 and 96% U-238 since the U-234 component is negligible.

$$\begin{aligned}
 \frac{1}{M_{\text{U}}} &= \frac{\omega_{\text{U-235}}}{M_{\text{U-235}}} + \frac{\omega_{\text{U-238}}}{M_{\text{U-238}}} = \frac{0.04}{235.04} + \frac{0.96}{238.05} \\
 M_{\text{U}} &= 237.9 \text{ g/mol}
 \end{aligned}$$

Next, use Equation (9) and the fact that $\rho_{\text{U}} = \rho_{\text{UO}_2} \frac{M_{\text{U}}}{M_{\text{UO}_2}}$

$$\begin{aligned}
 N_{\text{U-235}} &= \frac{\omega_{\text{U-235}} \rho_{\text{U}} N_{\text{Av}}}{M_{\text{U-235}}} = \frac{\omega_{\text{U-235}} N_{\text{Av}}}{M_{\text{U-235}}} \rho_{\text{UO}_2} \frac{M_{\text{U}}}{M_{\text{UO}_2}} \\
 &= \left(0.04 \frac{\text{g-}^{235}\text{U}}{\text{g - U}} \right) \frac{(6.022 \times 10^{23} \text{ atoms/mol})(10.5 \text{ g - UO}_2/\text{cm}^3)}{235.04 \text{ g-}^{235}\text{U/mol}} \left(\frac{237.9 \text{ g - U}}{[237.9 + 2(16.00)] \text{ g - UO}_2} \right) \\
 &= 9.49 \times 10^{20} \text{ atoms/cm}^3
 \end{aligned}$$